

Insecticide Guidelines

Richard G. St-Pierre, Ph.D. (January 2006)

Overview Of Insecticides Registered For Use In Saskatoon Orchards		
Insecticide	Registered Use	Timing
Bartlett Superior '70' Oil	Saskatoon bud moth, fruit tree leafrollers	Apply once per season in the spring at bud break
Orthene	Woolly elm aphid in newly-established, non-bearing saskatoon orchards, less than 4 years of age	Apply once per year in mid-July or early-August
Decis 5.0EC	Apple curculio, hawthorn weevil, lygus bug, saskatoon bud moth, saskatoon sawfly, tarnished plant bug	Apply 3 times per season: once at green tip, once at early flowering, once after petal drop but not within 21 days of harvest

Caution - This information is solely meant as a guide. Application of all pesticides must be in accordance with instructions on the product label as prescribed under the Pest Control Products Act. Always refer to the label.

Guidelines To The Use Of Bartlett Superior '70' Oil

Manufacturer - N.M. Bartlett Inc., 4509 Bartlett Rd., Beamsville, ON, L0R 1B1; 1 800 263 1287; www.bartlett.ca; 20 litres costs \$69.00 plus tax and shipping; 200 litres costs \$282.00 plus tax and shipping.

Trade Names - Bartlett Superior '70' Oil.

Formulation - 98.5% Superior '70' Oil.

Registered Use - Control of saskatoon bud moth and fruit tree leafrollers.

Timing Of Application - Apply once per season in the spring at bud break; Refer to Plate 1 and the Stages of Flower Bud Development color graphic in Section III: Botany of this manual for further information on timing.

Application Rate & Guidelines - Apply at the rate of 20 L product in 1000 L water per hectare. All plant surfaces are to be thoroughly coated.

Applicators should wear a long-sleeved shirt, long pants, chemical-resistant gloves, eye protection, and a mask fitted to exclude spray mist. Superior '70' Oil is an Eye Irritant.

Restrictions - Applications should not be made during freezing weather, or if frost is expected before the plants dry. Only one application per year is approved.

Points To Note - Horticultural spray oils such as Bartlett's Superior 70 Oil are highly-refined petroleum oils. If applied when plant buds are dormant, or shortly thereafter, such oils are effective against exposed eggs and soft-bodied immature and adult pests (including most species of mites and some species of young caterpillars and various sucking insects; they are also effective on eggs of mites, aphids, and some moths). Their mode of action is not clear but may include smothering of insect or eggs; they may also disrupt cell membranes. Modern horticultural oils are easy to use in the field, and are inexpensive and safe to handle. If properly used, they have low toxicity to most plants. They are relatively harmless to humans, other mammals and birds. Additionally, they dissipate quickly after spraying, leaving little or no residue on crops. Because of this reduced residual activity, they usually are less harmful to beneficial insects and predatory mites. Target pests develop little or no resistance to oils. The use of Bartlett's Superior 70 Oil as a replacement for at least the first application of Decis will help reduce the probability of insect pests developing resistance to Decis. Bartlett's Superior '70' Oil is well suited for use in Integrated Pest Management (IPM) programs, and can be used by organic growers.

Guidelines To The Use Of Orthene

Manufacturer - United Agri Products.

Trade Names - Orthene.

Formulation - Soluble powder; 75% acephate.

Registered Use - Control of woolly elm aphid in newly-established, non-bearing saskatoons, less than 4 years of age.

Timing Of Application - Apply once per year in mid-July or early-August.

Application Rate & Guidelines - Mix 1.7 gm / 2 L water (3.4 kg/ha with a planting density of 2,000 plants/ha). Apply 2 L per plant. Inject with a probe 15 cm from the stem of the plant to a depth of 12 cm at 3 to 5 locations around the plant. Inject into moist soil, otherwise a 2 L volume of insecticide may be insufficient to wet the volume of soil required for aphid control.

Restrictions - Use only on non-bearing saskatoons less than 4 years of age.

Points To Note - The efficacy of orthene tends to be greater if the application is made in early to mid-July. However, orthene is not registered for application at this time of the season.

Orthene is available in 1.5 kg bags; this is sufficient for approximately 880 plants at a cost of \$0.10 per plant.

Plans for a do-it-yourself single-point soil injector can be obtained from the PFRA Shelterbelt Center, Indian Head, SK.

Guidelines To The Use Of Decis 5.0 EC

Manufacturer - AgrEvo Canada Inc.

Trade Names - Decis 5.0 EC.

Formulation - 50 gm/L deltamethrin (synthetic pyrethroid); emulsifiable concentrate.

Registered Use - Control of apple curculio, hawthorn weevil, lygus bug, saskatoon bud moth, saskatoon sawfly, tarnished plant bug.

Timing Of Application - Apply 3 times per season; once at the green tip stage of flower bud development, once during early flowering (25 - 50% bloom), and once after petal drop, remembering to observe the preharvest interval of 21 days. Refer to Plate 1 and the Stages of Flower Bud Development color graphic in Section III: Botany of this manual for further information on timing.

Application Rate & Guidelines - Use 2 ml/4.5 L water for every 100 sq m of orchard row face (including both sides of the row alley, or both sides of a single row of plants; to calculate the total amount of Decis required, refer to the section on page 2 of this factsheet). Ensure that the foliage is thoroughly sprayed, including the undersides of the leaves.

Restrictions - Do not apply within 21 days of harvest. Do not apply when temperature is greater than 25°C; application in early-morning or early-evening is best.

Points To Note - Extensive use of Decis

may contribute to increasing problems with the McDaniel spider mite.

Decis controls insects through contact and/or ingestion. Decis works best under cool conditions. The first application will control the saskatoon bud moth, lygus and tarnished plant bug. The second application will control the saskatoon sawfly. The third application will ensure control of the saskatoon sawfly, hawthorn weevil and apple curculio.

Calculating Amount Of Pesticide

Required - 1) Multiply row length in m by row height in m
2) Multiply this figure by 2, then divide this last figure by 100
3) The result is the number of 100 square meter sections that must be sprayed
4) Multiply this figure by 2 ml to obtain the total amount of Decis required, and by 4.5 L to obtain the total amount of water required.
5) Calibrate sprayer and tractor speed to uniformly deposit spray volume over entire orchard area.

*Copyright 2006 by Richard G. St-Pierre, Ph.D.
www.prairie-elements.ca. All rights reserved. Any copying or publication or use of this publication or parts thereof for financial gain is not permitted. Users of this publication are allowed to print one (1) copy for personal use only. Otherwise, this publication may not be reproduced in any form, or by any means, in whole or in part for any purposes without prior written permission of the author. Due recognition must be given to the author for any use which may be made of any material in this publication. Requests for permission to copy or to make use of material in this publication, in whole or in part, should be addressed to: Richard St-Pierre, Email: prairie.elements@sasktel.net*

Disclaimer: This publication was designed to be an educational resource for individuals who are interested in

growing saskatoons, in orchards, shelterbelts, or gardens. Every effort has been made to ensure the accuracy and effectiveness of the information in this publication. However, the author makes no guarantee, express or implied, as to the information and procedures contained herein. The information cannot be guaranteed because knowledge of the biology and culture of the saskatoon may not be applicable to all locations every year. Additionally, the information that is available often changes over time. Little scientific research has been done on many aspects of the culture and management of saskatoons. Consequently, this publication can only serve as a guide. All actions taken which are based on the information presented in this publication are solely the responsibilities of the readers or users, and the author is not liable for any direct, indirect, incidental, or consequential damages in connection with or arising from the furnishing, performance, or use of this material. Comments on information contained in this publication are welcomed.